

SUPPLEMENTAL USE DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.

Please read the complete label affixed to the container for Discipline 2EC before applying. All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Use of Discipline 2EC according to this supplemental labeling is subject to the use precautions and limitations imposed by the label on the container.

IMPORTANT: Read the entire Directions for Use and the Limited Warranty and Disclaimer on the EPA-registered label for Discipline 2EC before using this product.

SOYBEANS (Not for use in California)		
PEST	Rate of Application	
	LB/AI/A	FL OZ/A
Alfalfa Caterpillar	0.033 to 0.10	2.1 to 6.4
Aphids		
Aster Leafhopper		
Bean Leaf Beetle		
Beet Armyworm*		
Cloverworm		
Corn Earworm		
Corn Rootworm Adult		
Cucumber Beetles		
Cutworms		
European Corn Borer		
Fall Armyworm		
Flea Beetle		
Grasshoppers		
Imported Cabbageworm		
Japanese Beetle Adult		
Leafhoppers		
Leafminer		
Loopers		
Mexican Bean Beetle Adult		
Pea Leaf Weevil		
Pea Weevil		
Plant Bug		
Saltmarsh Caterpillar		
Sap Beetle		
Southern Armyworm		
Stink Bugs		
Tarnished Plant Bug		
Thrips		
Tobacco Budworm*		
Webworms		
Western Bean Cutworm		
Whitefly		
Yellowstriped Armyworm		

SOYBEANS (Not for use in California)		
Lygus spp. Whitefly Two Spotted Spider Mite	0.08 to 0.10	5.12 to 6.4
REMARKS Apply in a minimum of 10 gallons per acre with ground equipment or 2 gallons per acre by aircraft at a minimum of 30 day intervals. Do not apply more than 0.3 lb ai per acre per season. Do not apply within 18 days of harvest. *Pyrethroid resistance is common for Beet Armyworm and Tobacco Budworm. Please consult your local or state agricultural authority to determine if resistance pest populations are in your area. If so refer to the resistance management statement in the DIRECTION FOR USE section of this label.		

TUBEROUS AND CORM VEGETABLES (Not for use in California)			
CROP	PEST	Rate of Application	
		LB/AI/A	FL OZ/A
Potato	Corn Wireworm	0.15 to 0.30	9.6 to 19.2
Sweet Potato	Tobacco Wireworm	(at-plant)	(at-plant)
Arracacha	Southern Potato		
Arrowroot	Wireworm	0.05 to 0.15	3.2 to 9.6
Chinese Artichoke	Japanese Beetle Grubs	(at cultivation	(at cultivation
Jerusalem Artichoke	June Beetle	or lay-by)	or lay-by)
Edible Canna	Sweetpotato Flea Beetle		
Cassava (Bitter and Sweet)	Cucumber Beetle	0.033 to 0.10	2.1 to 6.4
Chayobs (Root)	Sweetpotato Weevil	(foliar)	(foliar)
Chufa	Banded Cucumber Beetle		
Dasheen (Taro)	Black Flea Beetle		
Ginger	Whitefringed Beetle		
Laren	White Grub		
Tanier Turner	Sugarcane Beetle		
Yam Bean	Rootworms		
True Yam			
REMARKS Discipline 2EC may be applied as a soil incorporated broadcast, directed bed spray or a T-band spray into the planting furrow for the control of wireworms, rootworms, sweet potato flea beetle and white grubs. Apply Discipline 2EC at the rate of 0.15 to 0.3 pounds active (9.6 to 19.2 ounces formulated) per acre in a minimum of 10 gallons finished spray per acre. Discipline 2EC may be applied as one or more soil directed and incorporated treatments at cultivation or lay-by for the control of wireworms, rootworms and white grubs. Apply Discipline 2EC to the drill area and incorporate by cultivation equipment set to throw soil towards the drill area. Apply Discipline 2EC at a rate of 0.05 to 0.15 pounds active (3.2 to 9.6 ounces formulated) per acre in a minimum of 10 gallons finished spray per acre. Discipline 2EC may be applied as a foliar spray for the control of the adult life stages of flea beetles, click beetles (wireworms), cucumber beetles (rootworms), white fringed beetles and May/June beetles (white grubs). Apply Discipline 2EC at the rate of 0.033 to 0.1 lbs active (2.1 to 6.4 ounces formulated) per acre in a minimum of 10 gallons per acre by ground and 3 gallons per acre by air. Do not make more than 2 foliar applications per season no sooner than 21 days apart. Do not apply more than 0.5 lb. active ingredient per acre per season, including soil application. Do not apply within 21 days of harvest.			

Discipline is a registered trademark of AMVAC Chemical Corporation.

Registrant: AMVAC Chemical Corporation
 4100 East Washington Boulevard
 Los Angeles, CA 90023